

Label-free Chemical Imaging of Fungal Spore Walls by Raman Microscopy and Multivariate Curve Resolution Analysis

Hemanth Noothalapati^{1,*}, Takahiro Sasaki¹, Tomohiro Kaino^{1,2}, Makoto Kawamukai^{1,2},
Masahiro Ando³, Hiro-o Hamaguchi^{3,4} and Tatsuyuki Yamamoto^{1,2,*}

¹Raman Center for Medical and Biological Applications, Shimane University, Matsue 690-8504, Japan; ²Faculty of Life and Environmental Science, Shimane University, Matsue 690-8504, Japan; ³Consolidated Research Institute for Advanced Science and Medical Care, Waseda University, Tokyo 162-0041, Japan; ⁴ Institute of Molecular Science and Department of Applied Chemistry, National Chiao Tung University, Hsinchu 30010, Taiwan

* To whom correspondence may be addressed.

Dr. Hemanth Noothalapati

Raman Center for Medical and Biological Applications

Shimane University

1060, Nishikawatsu Cho, Matsue 690-8504, Shimane, Japan

Phone: +81-852-32-6571

Email: nvhnag@life.shimane-u.ac.jp

Or

Prof. Tatsuyuki Yamamoto


Faculty of Life and Environmental Science

Shimane University


1060, Nishikawatsu Cho, Matsue 690-8504, Shimane, Japan

Phone: +81-852-32-6551


Email: tyamamot@life.shimane-u.ac.jp





Supplementary Figure 1: Optical layout of Raman microscope used.



Supplementary Figure 2: MCR analyzed Raman spectra from **A)** 7 components model and **B)** 9 components model.



Supplementary Figure 3: Raman images constructed from MCR analysis from last three components shown in the main text. **A)** *S. pombe* vegetative cells and **B)** ascospores.



Supplementary Figure 4: 8 components AHCA. **A)** Dendrogram and **B)** Cluster spectra.